

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 29 November 2016		2. REPORT TYPE Briefing Charts		3. DATES COVERED (From - To) 02 November 2016 – 30 November 2016	
4. TITLE AND SUBTITLE A Hybrid Model for Multiscale Laser Plasma Simulations with Detailed Collisional Physics			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) David Bilyeu, Carl Lederman, Richard Abrantes, Hai Le			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER Q02Z		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQRS 1 Ara Drive Edwards AFB, CA 93524-7013			8. PERFORMING ORGANIZATION REPORT NO.		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQR 5 Pollux Drive Edwards AFB, CA 93524-7048			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-RQ-ED-VG-2016-328		
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited. The U.S. Government is joint author of the work and has the right to use, modify, reproduce, release, perform, display, or disclose the work.					
13. SUPPLEMENTARY NOTES For presentation at AFOSR Plasma and Electroenergetics Review Meeting; Arlington, VA (29 November 2016) PA Case Number: #16531; Clearance Date: 07 November 2016 Prepared in collaboration with ERC					
14. ABSTRACT Viewgraph/Briefing Charts					
15. SUBJECT TERMS N/A					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 23	19a. NAME OF RESPONSIBLE PERSON D. Bilyeu
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NO (include area code) N/A



A Hybrid Model for Multiscale Laser Plasma Simulations with Detailed Collisional Physics

*AFOSR Plasma and Electroenergetics Review Meeting
29-30 November 2016*



Collaborators:

- R. Caflisch (UCLA)
- A. Karagozian (UCLA)

Richard Abrantes^{1,2}, Hai Le^{1,2},
Carl Lederman²,
PI: David Bilyeu³
¹UCLA
²ERC Inc.
³AFRL/RQRS



Outline

- **Goals**
- **Review of Past Work**
- **Argon Collisional-Radiative Complexity Reduction Validation**
- **Non-Maxwellian CR**
- **Phase-accurate Multiscale Particle Push**





Goals

- Utilize hybridization techniques to produce accurate and efficient plasma simulations that spans many orders of magnitude in both space and time.
- Capture complex physics: excitation/ionization, transport, radiation, etc.
- Consistent collision operator across different levels of fidelity.

Current Focus:

- Generalization of collisional-radiative kinetics with level grouping
- General Hybridization techniques
- Focus on each solver before hybridization
- Special attention to low density low energy conditions

Laser Plasma Interaction



FRC



RQRS M&S Group



- **Government**

- Dr. David Bilyeu (LPI, in-space Chem)
- Dr. Justin Koo (Flight Support, Group Lead)
- Dr. Rob Martin (FRC)

- **Onsite Contractors (ERC inc.)**

- Richard Abrantes (Grad Student, LPI)
- Dr. Jun Araki (Flight Support, PIC)
- Dr. Carl Lederman (LPI)
- Dr. Michelle Scharfe (Flight Support)
- Dr. Eder Sousa (FRC)
- Jonathan Tran (Grad Student, Implicit PIC)

- **Summer Students**

- Astrid Raisanen
 - PhD UofM; Vlasov HET
- Daniel Crews
 - M.S. Washington; Collisionless Shock for V&V
- Kari Kawashima
 - B.S. UCLA; SM/MURF beta tester

- **Previous**

- Dr. Hai Le (LPI; now at Livermore)
- Dr. Artan Qerushi (FRC now at Lockheed)



Summary of Past Work

Maxwellian Inelastic Collisions


- Detailed CR model for multiple ionization stages
- Validation against experimental data
- Nonequilibrium radiation transport: coupling with a collisional-radiative model
- Inelastic collisions in a MF plasma: enhanced thermochemical kinetics.

Multiscale Hybridization

- A time-parallel/multiscale method with energy preservation

Analytical BGK

- Conservative even for large mass ratios
- Conservation are independent of collisional frequencies





Collisional Radiative (CR) Overview

Updates

- Expanded complexity reduction to include multiple ionization levels
- Adaptive integration technique -> fixes rate calculation for higher electron temperature
- Investigate grouping sensitivity
- Linked with LANL database for Argon cross sections and atomic level information
- Algorithms not hard coded for Argon.

Levels of Complexity

- Full rates (LANL)
- Cutoff nearly ionized levels
- Grouping Strategies
 - Uniform
 - Boltzmann
 - QSS (Boltzmann and Planck equilibrium)
- Group Selection
 - Electron configuration (no splitting information)
 - Highly excited states
 - Analysis of full run
 - Numerical optimization



CR Governing Equations

$$\begin{aligned} \left(\frac{df_e(\varepsilon, t)}{dt} \right)_{\text{coll, ex/dex}} = & - \sum_{m>n} N_n(f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(m|n)}^{e,\uparrow}(\varepsilon_0) \\ & + \sum_{m>n} N_m(f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \sigma_{(n|m)}^{e,\downarrow}(\varepsilon_1) \\ & + \sum_{m<n} N_m(f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \sigma_{(n|m)}^{e,\uparrow}(\varepsilon_1) \\ & - \sum_{m>n} N_n(f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(m|n)}^{e,\downarrow}(\varepsilon_0) \end{aligned}$$

$$\begin{aligned} \left(\frac{df_e(\varepsilon, t)}{dt} \right)_{\text{coll, i/r}} = & - \sum_n N_n(f_e(\varepsilon)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(+|n)}^{e,\uparrow}(\varepsilon_0) \\ & + N_+ \int_{\varepsilon_1} \int_{\varepsilon_2} (f_e(\varepsilon_1))(f_e(\varepsilon_2)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \left(\sqrt{\frac{2\varepsilon_2}{m_e}} \right) \left(\frac{\partial^2}{\partial \varepsilon_1 \partial \varepsilon_2} \sigma_{(n|+)}^{e,\downarrow}(\varepsilon_1, \varepsilon_2; \varepsilon_0) \right) d\varepsilon_1 d\varepsilon_2 \\ & - N_+(f_e(\varepsilon_2)) \left(\sqrt{\frac{2\varepsilon_2}{m_e}} \right) \int_{\varepsilon_0} \int_{\varepsilon_1} (f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \left(\frac{\partial^2}{\partial \varepsilon_1 \partial \varepsilon_2} \sigma_{(n|+)}^{e,\downarrow}(\varepsilon_1, \varepsilon_2; \varepsilon_0) \right) d\varepsilon_1 d\varepsilon_0 \\ & + \sum_n N_n \int_{\varepsilon_1} \int_{\varepsilon_0} \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) (f_e(\varepsilon_0)) \left(\frac{\partial^2}{\partial \varepsilon_1 \partial \varepsilon_2} \sigma_{(+|n)}^{e,\uparrow}(\varepsilon_2; \varepsilon_0, \varepsilon_1) \right) d\varepsilon_0 d\varepsilon_1 \end{aligned}$$

$$\begin{aligned} \left(\frac{dN_n(t)}{dt} \right)_{\text{coll, ex/dex}} = & - \sum_{m>n} N_n \int_{\varepsilon_0} (f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(m|n)}^{e,\uparrow}(\varepsilon_0) d\varepsilon_0 \\ & + \sum_{m>n} N_m \int_{\varepsilon_1} (f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \sigma_{(n|m)}^{e,\downarrow}(\varepsilon_1) d\varepsilon_1 \\ & + \sum_{m<n} N_m \int_{\varepsilon_1} (f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \sigma_{(n|m)}^{e,\uparrow}(\varepsilon_1) d\varepsilon_1 \\ & - \sum_{m<n} N_n \int_{\varepsilon_0} (f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(m|n)}^{e,\downarrow}(\varepsilon_0) d\varepsilon_0 \end{aligned}$$

$$\begin{aligned} \left(\frac{dN_n(t)}{dt} \right)_{\text{coll, i/r}} = & - N_n \int_{\varepsilon_0} (f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(+|n)}^{e,\uparrow}(\varepsilon_0) d\varepsilon_0 \\ & + N_+ \int_{\varepsilon_0} \int_{\varepsilon_1} \int_{\varepsilon_2} (f_e(\varepsilon_1))(f_e(\varepsilon_2)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \left(\sqrt{\frac{2\varepsilon_2}{m_e}} \right) \left(\frac{\partial^2}{\partial \varepsilon_1 \partial \varepsilon_2} \sigma_{(n|+)}^{e,\downarrow}(\varepsilon_1, \varepsilon_2; \varepsilon_0) \right) d\varepsilon_1 d\varepsilon_2 \end{aligned}$$



CR Governing Equations cont.

- Current Assumptions**

- Electron dominated collisions
- δ function for ion distribution
- Maxwellian electrons

- Current Model Includes**

- Multiple ionization levels
- Excitation/de-excitation
- Ionization/recombination
- Multi-photon ionization and inverse Bremsstrahlung
- Radiation losses via, Bound-Bound and Bound-Free

*previous included

$$\begin{aligned} \frac{dn_n^{+k}}{dt} = & - \sum_{m>n} \alpha_{(m|n)}^{+k,e} N_e N_{+k,n} + \sum_{m>n} \beta_{(n|m)}^{+k,e} N_e N_{+k,m} + \sum_{m>n} A_{(n|m)}^{+k} N_{+k,m} \\ & + \sum_{m<n} \alpha_{(n|m)}^{+k,e} N_e N_{+k,m} - \sum_{m<n} \beta_{(m|n)}^{+k,e} N_e N_{+k,n} - \sum_{m<n} A_{(m|n)}^{+k} N_{+k,n} \\ & - \sum_j \alpha_{(+,j|n)}^{+k,e} N_e N_{+k,n} + \sum_j \beta_{(n|+,j)}^{+k,e} N_{+(k+1),j} N_e^2 \\ & \alpha_{(m|n)}^e = \int_{E_{nm}}^{\infty} \sigma_{nm}^e(\varepsilon) v_e f(v_e) dv_e \\ & \beta_{(n|m)}^e = \frac{n^2}{m^2} e^{+x_{nm}} \alpha_{(m|n)} \quad A_{(n|m)} = \left(\frac{8\pi^2 e^2}{m_e c^3} \right) \frac{g_n}{g_m} f_{nm} \end{aligned}$$



CR Grouping Techniques

Uniform

- Solely conserves number density
- Weights each level according to level degeneracy within group

Conserved Variable:


$$\tilde{N}_n = \sum_{i \in n} N_i$$

Effective rate:

$$\tilde{\alpha}(m|n) = \sum_{i \in n} \frac{g_i}{g_n} \sum_{j \in m} \alpha(j|i)$$

$$\frac{dN_n}{dt} = N_e \left[\sum_{m>n} \alpha_{(m|n)} N_n + \sum_{m<n} \beta_{(m|n)} N_n \right] + \dots$$

$$\frac{d\tilde{N}_n}{dt} = N_e \tilde{N}_n \left[\sum_{m>n} \tilde{\alpha}_{(m|n)} + \sum_{m<n} \tilde{\beta}_{(m|n)} \right] + \dots$$



Boltzmann

- Conserves number density
- Preserves energy in groups through group temperature description

Conserved Variable:

$$N_{n_0} \& N'_n = \frac{N_{n_0}}{g_{n_0}} \sum_{i \in n} g_i e^{-\Delta E_i / T_n}$$

Effective rate:

$$\begin{aligned} & \tilde{\alpha}(m'|n') \\ &= \sum_{i \in n} \frac{g_i e^{-\Delta E_i / T_n}}{Z'_n} \sum_{j \in m'} \alpha(j|i) \end{aligned}$$



Complexity Reduction for Argon

Full Lines (LANL)

- Based on quantum calculations with corrections for low temperature

NIST Cutoff


- Starts with LANL and assumes higher excited states are ionized
- Cutoff experimentally determined
- 2-3x reduction

Electron Configuration

- Groups based on electron configuration
- Uses uniform grouping
- 10-15x reduction over NIST

Grouping

- Boltzmann or Uniform grouping
- Saves 20-30% over Electron Splitting
- Case by case basis





Argon Level Grouping Isothermal Test Setup



Simulation Setup

- Pressure: 4.22 Torr (5.55×10^{-3} atm)
- Ion Temperature: 0.035 eV
- Atomic Density: 10^{20} 1/m³
- Ionization fraction: 10^{-13}
- Electron Temperature: 10 & 100 eV
- $t = [0, 10^6]$ seconds


Groupings

- NIST cutoff with electron grouping
- NIST cutoff with electron grouping and Boltzmann grouping
- NIST cutoff with electron grouping and Uniform grouping



Argon Level Grouping Isothermal Test

Electron Temperature 10 eV




Argon +1
12 electron configurations
9 Isolates states
1 group with 3 states

Uniform group propagates
errors to isolated states



Argon Level Grouping Isothermal Test

Electron Temperature 100 eV




Argon +3
17 electron configurations
14 Isolates states
1 group with 3 states

Uniform groups propagates
errors to isolated states



Argon Level Grouping Isothermal Test Error



ODE Solver tolerance
Relative error 1^{-4}
Absolute error 1
~1200 Radau5 time steps
dt increases exponentially

Solution improves if
either tolerance is
decreased but at the
expense of computational
time. E.g. relative error
 10^{-6} -> computational
time triples



Non-Maxwellian Electrons CR - Ions (Preliminary)



• Current Assumptions

- Electron dominated collisions
- Single ionization level
- Isotropic EEDF
- δ function for ion distribution

$$\left(\frac{dN_n(t)}{dt}\right)_{\text{coll,ex/dex}} = - \sum_{m>n} N_n \int_{\varepsilon_0} (f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}}\right) \sigma_{(m|n)}^{e,\uparrow}(\varepsilon_0) d\varepsilon_0$$

$$+ \sum_{m>n} N_m \int_{\varepsilon_1} (f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}}\right) \sigma_{(n|m)}^{e,\downarrow}(\varepsilon_1) d\varepsilon_1$$

$$+ \sum_{m<n} N_m \int_{\varepsilon_1} (f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}}\right) \sigma_{(n|m)}^{e,\uparrow}(\varepsilon_1) d\varepsilon_1$$

$$- \sum_{m<n} N_n \int_{\varepsilon_0} (f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}}\right) \sigma_{(m|n)}^{e,\downarrow}(\varepsilon_0) d\varepsilon_0$$

• Current Model Includes

- Elastic electron collisions
- Excitation/de-excitation
- Ionization/recombination

$$\left(\frac{dN_n(t)}{dt}\right)_{\text{coll,i/r}} =$$

$$- N_n \int_{\varepsilon_0} (f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}}\right) \sigma_{(+|n)}^{e,\uparrow}(\varepsilon_0) d\varepsilon_0$$

$$+ N_+ \int_{\varepsilon_0} \int_{\varepsilon_1} \int_{\varepsilon_2} (f_e(\varepsilon_1))(f_e(\varepsilon_2)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}}\right) \left(\sqrt{\frac{2\varepsilon_2}{m_e}}\right) \left(\frac{\partial^2}{\partial \varepsilon_1 \partial \varepsilon_2} \sigma_{(n|+)}^{e,\downarrow}(\varepsilon_1, \varepsilon_2; \varepsilon_0)\right) d\varepsilon_1 d\varepsilon_2$$



Non-Maxwellian Electrons CR - Electrons (Preliminary)



• Current Assumptions

- Electron dominated collisions
- Single ionization level
- Isotropic EEDF
- δ function for ion distribution

$$\begin{aligned} \left(\frac{df_e(\varepsilon, t)}{dt} \right)_{\text{coll, ex/dex}} = & - \sum_{m>n} N_n(f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(m|n)}^{e,\uparrow}(\varepsilon_0) \\ & + \sum_{m>n} N_m(f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \sigma_{(n|m)}^{e,\downarrow}(\varepsilon_1) \\ & + \sum_{m<n} N_m(f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \sigma_{(n|m)}^{e,\uparrow}(\varepsilon_1) \\ & - \sum_{m>n} N_n(f_e(\varepsilon_0)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(m|n)}^{e,\downarrow}(\varepsilon_0) \end{aligned}$$

• Current Model Includes

- Elastic electron collisions
- Excitation/de-excitation
- Ionization/recombination


$$\begin{aligned} & \left(\frac{df_e(\varepsilon, t)}{dt} \right)_{\text{coll, i/r}} = \\ & - \sum_n N_n(f_e(\varepsilon)) \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) \sigma_{(+|n)}^{e,\uparrow}(\varepsilon_0) \\ & + N_+ \int_{\varepsilon_1} \int_{\varepsilon_2} (f_e(\varepsilon_1))(f_e(\varepsilon_2)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \left(\sqrt{\frac{2\varepsilon_2}{m_e}} \right) \left(\frac{\partial^2}{\partial \varepsilon_1 \partial \varepsilon_2} \sigma_{(n|+)}^{e,\downarrow}(\varepsilon_1, \varepsilon_2; \varepsilon_0) \right) d\varepsilon_1 d\varepsilon_2 \\ & - N_+(f_e(\varepsilon_2)) \left(\sqrt{\frac{2\varepsilon_2}{m_e}} \right) \int_{\varepsilon_0} \int_{\varepsilon_1} (f_e(\varepsilon_1)) \left(\sqrt{\frac{2\varepsilon_1}{m_e}} \right) \left(\frac{\partial^2}{\partial \varepsilon_1 \partial \varepsilon_2} \sigma_{(n|+)}^{e,\downarrow}(\varepsilon_1, \varepsilon_2; \varepsilon_0) \right) d\varepsilon_1 d\varepsilon_0 \\ & + \sum_n N_n \int_{\varepsilon_1} \int_{\varepsilon_0} \left(\sqrt{\frac{2\varepsilon_0}{m_e}} \right) (f_e(\varepsilon_0)) \left(\frac{\partial^2}{\partial \varepsilon_1 \partial \varepsilon_2} \sigma_{(+|n)}^{e,\uparrow}(\varepsilon_2; \varepsilon_0, \varepsilon_1) \right) d\varepsilon_0 d\varepsilon_1 \end{aligned}$$

$$\left(\frac{df_e(\varepsilon, t)}{dt} \right)_{\text{elastic}} = v_{ee}(F_e - f_e) + v_{ei}(F_{ei} - f_e)$$



Non-Maxwellian CR Numerical Challenges

- Long complex CR formulas
 - Stiff equations
 - Range of scales
 - Boundary conditions
 - Multi-dimensional integrations
-
- Hydrogen recombination example:
 - $e_0 + H_n \leftarrow e_1 + e_2 + H_+$
 - Energy equation in terms of the electron's kinetic energies, ε , and ionization energy I_n .
 - $\varepsilon_0 = \varepsilon_1 + \varepsilon_2 + I_n$
 - Evaluating the effect of recombination on a single species of hydrogen requires the evaluation of a 2D integral.





Hybridization Techniques

- Solve ODE/PDE with a coarse “C” solver and fine “F” solver
- Coarse propagates less information and is more computationally efficient
- Fine propagates more information and is more accurate
- User defined coarse error function $h(u), 0 \leq h \leq 1$
- when $h=0$ coarse is accurate
- Q compression operator
- R reconstruction operator
- $h, \epsilon, Q, \& R,$ are problem dependent

Simpler hybrid (H) method


$$H1_{[t+dt,t]} = \begin{cases} C_{[t+dt,t]}(Q(\text{if needed})) & \text{if } h \leq \epsilon \\ F_{[t+dt,t]}(R(\text{if needed})) & \text{if } h > \epsilon \end{cases}$$

More complex time parallel (TP) method

$$H2_{[Jdt,0]}(u(0)) = \begin{cases} C_{[Jdt,0]}Q(u(0)) & \text{if } h \leq \epsilon \\ C_{[Jdt,0]}Q(u(0)) + \left(\sum_{j=0}^{j=J-1} RC_{[Jdt,(j+1)dt]}QF_{(j+1)dt,jdt}RC_{[jdt,0]}Q(u(0)) \right) & \text{if } \epsilon < h \leq \gamma \\ \left(\sum_{j=0}^{j=J-1} RC_{[Jdt,(j+1)dt]}C_{(j+1)dt,jdt}C_{[jdt,0]}Q(u(0)) \right) & \\ F_{[Jdt,0]}(u(0)) & \text{if } h > \gamma \end{cases}$$



Hybridization Charged Particle in Magnetic Mirror



$$B_x = -\frac{2Cxz^3}{a^4}$$
$$B_y = -\frac{2Cyz^3}{a^4}$$
$$B_z = C\left(1 - \frac{z^4}{a^4}\right)$$

Gyrokinetic + Phase and Time Parallel method are the most efficient

Simple blending could be more accurate as the problem approaches steady state

*Submitted to Journal



Summary



- **Collisional Radiative**

- Boltzmann grouping improves group representation over applied uniform distribution
- Minimization of error, which is expected to accumulate quickly during highly-transient durations, and acceleration of method makes Boltzmann reduction a strong case for future coupled simulations
- Adaptive integration allowed for faster simulations at larger electron temperatures
- Sensitive to selected groups
- Grouping reduces stiffness
- Robust solver capable of handling, T_e from 1 – 1,000 eV and 10^{16} - 10^{24} particles
- Initial work on non-Maxwellian CR has begun and early numerical issues have been addressed

- **Hybridization**

- Shows that hybridization technique can be more computational efficient than the schemes that comprise it



Future Work

- **Collisional Radiative Simulations**
 - Further comparisons between reduced mechanisms with QSS
 - Line identification and width assignments in conjunction with experimental spectra
 - Apply Boltzmann grouping to new CR Argon rates and test with 1D MHD Argon shock by Kapper, et.al. Future plans to extend to 2 and 3D MF
 - Rerun previous LPI test case with level grouping; laser source term, heavy-electron elastic collisions, and multi-electron energy correction and heavy energy equation (Te, Th)
- **Group selection through numerical analysis, optimization**
- **Hybridize Maxwellian CR with QSS**
- **Non-Maxwellian finite rate EEDF implementation**



Questions?